


0277495... 888. 2

ISBN 82-7257-295-8



## **NPD-BULLETIN NO 5**

A revised Cretaceous and Tertiary  
lithostratigraphic nomenclature  
for the Norwegian  
North Sea

Edited by  
D. Isaksen and K. Tonstad

**Oljedirektoratet**  
Desember 1989

## CONTENTS

	Page		Page
INTRODUCTION .....	3	Jorsalfare Formation .....	30
Background .....	3	Hardråde Formation .....	30
Participation .....	3	Undifferentiated Shetland Group .....	32
Editorial statement .....	3		
LITHOSTRATIGRAPHIC PROCEDURES .....	3	TERTIARY .....	32
General .....	3	Basin evolution .....	32
Presentation of data .....	3	Lithostratigraphic notes .....	35
CHRONOSTRATIGRAPHIC FRAMEWORK .....	3	Rogaland Group .....	35
STRUCTURAL NOMENCLATURE .....	3	Hordaland Group .....	35
GEOLOGICAL INTRODUCTION .....	3	Nordland Group .....	35
CRETACEOUS .....	6	REVISED TERTIARY LITHOSTRATIGRAPHY .....	35
Basin evolution .....	6	Rogaland Group .....	35
Lithostratigraphic notes .....	7	Våle Formation .....	35
Cromer Knoll Group .....	7	Maureen Formation .....	35
Shetland Group .....	7	Ty Formation .....	39
REVISED CRETACEOUS		Vidar Formation .....	39
LITHOSTRATIGRAPHY .....	16	Lista Formation .....	40
Cromer Knoll Group .....	16	Andrew Formation .....	41
Åsgard Formation .....	17	Heimdal Formation .....	41
Tuxen Formation .....	17	Meile member .....	42
Mime Formation .....	19	Sele Formation .....	42
Sola Formation .....	19	Fiskebank Formation .....	44
Rødby Formation .....	20	Forties Formation .....	44
Agat Formation .....	21	Hermod Formation .....	45
Ran sandstone units .....	21	Balder Formation .....	46
Shetland Group .....	23	Hordaland Group .....	48
Hidra Formation .....	24	Frigg Formation .....	49
Hod Formation .....	24	Grid Formation .....	49
Tor Formation .....	25	Skade Formation .....	52
Ekofisk Formation .....	26	Vade Formation .....	54
Svarte Formation .....	27	Nordland Group .....	54
Blodøks Formation .....	28	Utsira Formation .....	54
Tryggvason Formation .....	28	LIST OF FIGURES .....	56
Kyrre Formation .....	29	REFERENCES .....	57

## INTRODUCTION

### Background

Deegan & Scull (1977) published a lithostratigraphic nomenclature for the central and northern North Sea. In early 1980 an initiative was taken by the exploration group of NIFO (the Norwegian Industrial Federation of Operating Companies) to consider a revision of the existing lithostratigraphy. The intention was to revise all the periods, but later it was decided to postpone work on the Cretaceous and Tertiary until the committees working on the Triassic and Jurassic had completed their task. The revised nomenclature for the Triassic and Jurassic of offshore Norway was published in 1984 (Vollset & Dore 1984).

In May 1987 the Norwegian Petroleum Directorate (NPD) proposed that the Cretaceous and Tertiary nomenclature should be revised. Den norske stats oljeselskap, Statoil, Norsk Hydro a.s. and Saga Petroleum a.s. were invited to participate in the work. The reason for inviting only these three operators was to keep the committees to a reasonable size.

### Participation

The working party was organised as two committees, both chaired by Dag Isaksen from NPD (Mobil Exploration from 1.9.88). Kjetil Tonstad from NPD joined the group when Dag Isaksen left NPD, and the groups were later chaired by Kjetil Tonstad and Dag Isaksen in cooperation.

#### *Cretaceous:*

Jon Einar Tellefsen (Esso Norge from 1.10.88) and Erik Holtar from Norsk Hydro, Per Audun Hole, Kjell Sigve Lervik and Thorbjørn Monsen from Statoil, Bjørn Thorleif Gunnar Wandås and Stein Nybakken from Saga Petroleum, Dag Isaksen and Kjetil Tonstad from NPD.

#### *Tertiary:*

Anne Strømme Lycke and Erik Holtar from Norsk Hydro, Kjell Sigve Lervik and Gro Kyllingstad from Statoil, Thomas Hardt from Saga Petroleum, Dag Isaksen and Kjetil Tonstad from NPD.

Valuable help and comments have been received from other geologists in Saga Petroleum, Norsk Hydro, Statoil and the Norwegian Committee on Stratigraphy (Norsk Stratigrafisk Komité). Foreign operators have commented on the text. In addition we have enjoyed valuable cooperation with the Geological Survey of Denmark (Danmarks Geologiske Undersøgelse) with regard to the Lower Cretaceous.

### Editorial statement

This paper proposes major changes to the lithostratigraphy in the Norwegian sector. However, some parts of the nomenclature proposed by Deegan & Scull (1977) remain unaltered or only modified, and the work of those authors has sometimes been almost directly quoted.

The introduction and general notes on the Cretaceous and Tertiary have been written by the editors on the basis of contributions from committee members. The authors of the formal descriptions are listed in alphabetical order at the beginning of each of the two main sections.

## LITHOSTRATIGRAPHIC PROCEDURES

### General

The lithostratigraphic nomenclature presented is based on the guidelines of the International Subcommission on Stratigraphic Nomenclature (Hedberg 1976) and "Regler og råd for navnsetting av geologiske enheter i Norge" (Nystuen 1986).

The criteria used for selecting type and reference wells are completeness of section, quality of wire-line logs and availability of cores. Reference wells are also used to illustrate different developments and boundaries of formations.

In this paper we have named most new formations from Norse mythology or after Norwegian "Viking" kings. Formations that are already formally defined retain their names. The overall nomenclature may thus therefore appear heterogeneous.

We have limited the nomenclature to two categories of lithostratigraphic units, namely group and formation. We have, however, found it necessary to suggest some additional informal subdivisions (see General lithostratigraphic notes).

### Presentation of data

The groups and formations are defined under the following headings: name, well type section, well reference section, thickness, lithology, basal stratotype, characteristics of the upper boundary, distribution, age and depositional environment.

The vertical scale on well logs in this report is 1:2000. Depths are given on all figures. All depths quoted are from RKB. All units defined in the paper are listed alphabetically at the front. The lithological legend is given in Fig. 1.

Illustrations also comprise distribution maps, seismic profiles and cross sections.

## CHRONOSTRATIGRAPHIC FRAMEWORK

Ages referred to in the text are at stage level for the Cretaceous and series level for the Tertiary (Fig. 2) Boreal terminology is adopted for the latest Jurassic - earliest Cretaceous stages, using the Volgian as the final Jurassic stage and the Ryazanian as the initial Cretaceous one.

## STRUCTURAL NOMENCLATURE

The structural features referred to in this paper are outlined on Fig. 3. Their names are suggested by a committee working on structural geology (Brekke et al., in prep.) to become formalised.

## GEOLOGICAL INTRODUCTION

To understand the Cretaceous and Tertiary development of the North Sea requires familiarity with the pre-Cretaceous geological history. The geological history of the North Sea is therefore briefly reviewed in this introduction. For more details see P.A. Ziegler (1981) and W. H. Ziegler et al. (1986).

### LITHOLOGICAL LEGEND

	Shale/Claystone		Lignite/Coal		Limestone streaks
	Siltstone		Carbonaceous		Dolomite streaks
	Sandstone		Bituminous		Unconformity
	Conglomerate		Chert		Pyrite
	Marl		Argillaceous		Glaucinite
	Limestone		Sand streaks		Fossils, unspecified
	Dolomite		Very sandy		
	Salt		Sandy		
	Gypsum		Slightly sandy		
	Anhydrite		Silty		
	Tuff		Silt streaks		

Fig. 1

### CRETACEOUS

U P P E R	MAASTRICHTIAN
	CAMPANIAN
	SANTONIAN
	CONIACIAN
	TURONIAN
	CENOMANIAN
L O W E R	ALBIAN
	APTIAN
	BARREMIAN
	HAUTERIVIAN
	VALANGINIAN
	RYAZANIAN

### TERTIARY/QUATERNARY

Quat	PLEISTOCENE
T E R T I A R Y	PLIOCENE
	MIOCENE
	OLIGOCENE
	EOCENE
	PALEOCENE
	Danian

Stage nomenclature for Cretaceous and series nomenclature for the Tertiary and the Quaternary.

Fig. 2.




Fig. 3

The Phanerozoic history can be divided into two major phases. The first phase is related to generation of the Pangaeon supercontinent. In the North Sea region the Caledonian (Ordovician-Devonian) and Hercynian orogenies (Carboniferous-Permian) produced the major events during this phase.

During the Caledonian orogeny the collision between the Fennoscandian Baltic Shield and the Laurentian-Greenland Shield led to formation of compressional structural elements that were superimposed on older Precambrian structural trends. The Caledonian structural elements are typically directed northeast-southwest. They were reactivated during subsequent tectonic events.

The second major tectonic phase was associated with the break-up of the Pangaeon supercontinent. In late Palaeozoic and early Triassic times tensional forces acted on the whole area and the supercontinent began to break up. The fragmentation culminated with the opening of the North Atlantic during the Tertiary. From Permian time onwards the area has been dominated by extension which produced grabens and other basins. Changes in sea level have been the main factor controlling the sedimentary processes.

A new and important structural feature developed in the North Sea during the Permian. The Mid North Sea High is a major east-west trending high dividing the North Sea area into a northern and a southern salt basin.

The Permian succession can generally be divided into a lower, sandy unit (Rotliegendes) and an upper unit (Zechstein) dominated by marine evaporites. These units can be recognised in the southern and central North Sea, but in the northern North Sea the nature of the Permian sediments is more uncertain. In the southern and central North Sea the salt sequences have been mobile since Triassic times, complicating the tectonic picture of the area.

In the Norwegian part of the North Sea the Triassic period was dominated by clastic sedimentation in a subsiding rift system. During the Jurassic and earliest Early Cretaceous the North Sea was affected by a series of earth movements in the Kimmerian tectonic episode. Rifting, with associated block faulting and erosion, took place in the later part of the Jurassic, and sedimentation was dominated by clastics.

## CRETACEOUS

### Basin evolution

A major phase of uplift and erosion producing the Late Kimmerian movements, took place across north-western Europe during late Volgian to Ryazanian time. This resulted in widespread regression which formed isolated sedimentary basins where deposition took place under dominantly anaerobic, reducing bottom conditions (Rawson & Riley 1982). In these basinal areas sedimentation was continuous from Late Jurassic to Early Cretaceous. These sediments belong to the Draupne, Mandal and Flekkefjord Formations (Vollset & Doré 1984).

The regression culminated in the Middle-Late Ryazanian and affected the North Sea.

The Early Cretaceous was essentially a period of transgression with minor regressions. (Rawson and

Riley 1982). The anaerobic conditions that dominated during the Late Volgian-Early Ryazanian regression ceased (Hesjedal & Hamar 1983). During the Valanginian to late Barremian times relatively quiet conditions dominated with sedimentation of shales and marls (Åsgard, Mime and Tuxen Formations, see Fig. 4).

During the transgression the sea covered progressively higher areas. Under these conditions condensed shallow-marine carbonates developed (Mime Formation). The present distribution of these limestones therefore reflects the subsidence pattern of the topographically higher features.

During Mid-Late Aptian times a regression occurred, which, together with movements along the North Sea rifts caused a change in lithology. In the basinal areas the claystones changed from calcareous rich to more organic rich (Sola Formation). Sandstones (Agat Formation and Ran sandstone units) were simultaneously deposited in some areas as submarine fans, due to erosion along the flanks of structural highs (Måløy Fault Blocks, Fladen Ground Spur/Andrew Ridge, Utsira High, Jæren High and Sele High).

The Mid-Late Aptian regressive event was followed by a regional transgression during Albian time. The sea flooded onto the structural highs, only the highest parts of which were exposed to erosion. Sandstone sequences (Agat Formation and Ran sandstone units) continued to be deposited along the flanks of structural highs as the sea encroached onto areas close to exposed Jurassic, Triassic or older sandy formations. The organic shales passed into a new fine-grained calcareous unit (Rødby Formation). The carbonate content continued to increase towards the end of the Albian.

A Late Albian-Early Cenomanian regressive event led to erosion and/or non-deposition of sediments along the flanks of structural highs. The Late Albian-Early Cenomanian sequences are very thin or absent in many wells.

The Upper Cretaceous sequence in the North Sea Basin was deposited in an open marine environment. In northwest Europe there was a general rise in relative sea level during the Late Cretaceous (Hancock & Kauffman 1979), although eustatic sea level on a global scale dropped overall during Turonian-Maastrichtian time (Haq et al. 1987). The Cretaceous subsidence in the Viking Graben and Central Trough has been related to lithospheric cooling after the Middle to Late Jurassic rifting phase (McKenzie 1978, Sclater & Christie 1980). This subsidence was increased by isostatic response to sediment loading (Beaumont & Sweeney 1978).

The Late Cretaceous was a quiet tectonic period. In the southern and central North Sea the supply of terrigenous material was reduced from the transition to the Cenomanian onwards, and pure carbonates were deposited (Shetland Group chalk facies).

In the northern North Sea the Late Cretaceous is dominated by a continuous, argillaceous and calcareous marine sequence (Shetland Group siliclastic facies).

It has been inferred that sedimentation rates somewhat exceeded the combined effects of subsidence and rising sea level in the Viking Graben (Ziegler 1982). Thus, water depth decreased gradually, though true shallow-water conditions were never established (Watts et al. 1980). During the Maastrichtian, the East

Shetland Basin and the Horda Platform were submerged and surrounding land areas comprised the Shetland Platform, Scotland, Norway and Greenland (Ziegler 1982, Hancock 1984).

#### General lithostratigraphic notes

The changes that have been made to the Cretaceous lithostratigraphy are documented in the group/formation definitions which follow. However, a brief summary of the principal revisions is given below.

#### *Cromer Knoll Group:*

The Valhall Formation, as defined by Deegan & Scull (1977), embraces all Lower Cretaceous sediments below the Rødby Formation in the Norwegian sector. The wealth of new data since the time of that publication allows several new formations to be separated from the Valhall Formation. Hesjedal & Hamar (1983) informally named some new formations (the Sola, Utvik, Klepp, Florø and Kopervik Formations) within the Valhall Formation. Jensen et al. (1986) formally defined two new formations (the Tuxen and Sola Formations), retaining the term Valhall Formation for the remnant of the original formation. In this report five formations are defined below the Rødby Formation: the Åsgard, Tuxen, Sola, Mime and Agat Formations. In addition the informal Ran sandstone units are described. An idealised development of the Cromer Knoll Group in the northern and central North Sea is shown in Fig. 4.

The Åsgard Formation consists of marly facies, and comprises the remainder of the original Valhall Formation after subdivision. According to Hedberg (1976), a previously established formation which is subdivided into new units must either be raised to group rank or abandoned. The old name should not be retained for any divisions of the original unit. Since the Cromer Knoll is well established as the group name for the Lower Cretaceous (also used on Haltenbanken by Dalland et al. 1988), we found it impractical to upgrade the Valhall Formation, and have therefore abandoned it.

The Sola Formation, which is part of the former Valhall Formation, was described and named, but not formally defined, by Hesjedal & Hamar (1983). Jensen et al. (1986) formally defined the formation. Even though the name conflicts with the rules laid down by Nystuen (1986), Sola being a village in Norway, it is retained in the Norwegian sector since the formation has already been defined in the Danish sector.

In their 1983 publication, Hesjedal & Hamar also informally named a basal limestone found in connection with structural highs, and called it the Utvik Formation. It is formally defined in this paper as the Mime Formation.

Sandstone intervals were developed in the vicinity of some structural highs at various times during the Early Cretaceous. Since it is uncertain how far they are related to each other we have found it expedient not to formally name individual sandstones. Their varying age also prohibits placing them as a member in one of the defined formations in the Cromer Knoll Group. We therefore propose to name them informally as the Ran sandstone units.

The Lower Cretaceous sandstones found in the Agat Field are thicker and can be correlated between wells.

We have therefore given these sandstones formation status, the Agat Formation.

The Tuxen Formation defined by Jensen et al. (1986) is extended into the Norwegian sector.

#### *Shetland Group:*

The Shetland Group, as introduced by Deegan & Scull (1977), included the Upper Cretaceous siliciclastic sequence in the northern North Sea. The Shetland Group, as defined herein, has been expanded to include both the original Shetland Group and the former Chalk Group (Deegan & Scull 1977). There are several reasons for combining these Upper Cretaceous groups. The relationship between the Shetland and Chalk Groups was not properly established by Deegan & Scull (1977), owing to the limited data available at that time.

There is good correlation between the formation boundaries within the siliciclastic facies of the former Shetland Group in the northern North Sea and the chalk facies of the former Chalk Group in the central North Sea. Their boundaries are in chronostratigraphic accordance and there are great similarities in log patterns in the two regions despite the differences in lithological facies (Fig. 5).

Interfingering of the various lithologies and the presence of transitional lithologies have created difficulties in the transition zone between the two facies. Hence, it has been found most convenient to separate the facies on formation level. The previous formation subdivision has therefore been retained in the new Shetland Group.

With the amalgamation of the former Chalk and Shetland Groups, the term Chalk Group was discarded for two reasons. According to the rules for naming lithostratigraphic units, lithological terms should be avoided as unit names. The term Shetland Group was therefore chosen for the Upper Cretaceous sequence because it was already established in the northern North Sea and on the central and northern Norwegian Shelf (Dalland et al. 1988).

In the chalk facies area, no major changes have been proposed in the nomenclature used by Deegan & Scull (1977) except that the name "Plenus Marl Formation" has been changed to "Blodøks Formation", this name being used in both the chalk and siliciclastic facies areas. The term Plenus Marl Formation was renamed to accord with the rules for naming lithostratigraphic units, since fossil names should be avoided for stratigraphic units.

In the siliciclastic facies area, the informal subdivision proposed by Deegan & Scull (1977) has been formalised by defining the formations. The names Svarte, Blodøks, Tryggvason, Kyrre and Jorsalfare Formations have been introduced for the former Formations A, B, C, D and E. Formation D is equivalent to the Flounder Formation which is the name of the siliciclastic part of the former Chalk Group in the central part of the central North Sea (Fig. 6).

A new name, Hardråde Formation, has been introduced for the unit representing the Maastrichtian interval in the Shetland Group on the Horda Platform. The lower part of the group in this area is not formally subdivided. Although present in the Troll area, this lower part of the succession has not been found in other wells on the Horda Platform. The development of the sequence here, also differs from that in the Viking

Idealised development of the Cromer Knoll Group in the Northern and Central North Sea.




Fig. 4 Development of the Cromer Knoll Group

R-0168






Fig. 5 Well correlation in the Central trough and the Viking Graben

SERIES	STAGE	Deegan & Scull (1977)		Present subdivision			
		Chalk Gp.	Shetland Gp.	Shetland Gp.			
		<sup>W</sup> Central North Sea <sup>E</sup>	<sup>W</sup> Northern North Sea <sup>E</sup>	<sup>W</sup> Central North Sea <sup>E</sup>	<sup>W</sup> Northern North Sea <sup>E</sup>		
Paleocene	Thanetian	Maureen Fm./ Unnamed unit		Lista / Våle / Maureen / Ty Fm.			
	Danian	Ekofisk Fm.	Fm. F	Ekofisk Fm.			
Upper Cretaceous	Maastrichtian	Tor Fm.		Fm. E	Tor Fm.	Jorsalfare Fm.   Hardråde Fm.	
	Campanian	Flounder Fm.	Hod Fm.	Fm. D	Flounder Fm.	Hod Fm.	Kyrre Fm.
	Santonian						
	Coniacian						
	Turonian	Herring Fm.		Fm. C	Herring Fm.	Tryggvason Fm.	Unspec.
		Plenus Marl Fm.		Fm. B	Blødøks Fm.		
	Cenomanian	Hidra Fm.		Fm. A	Hidra Fm.	Svarte Fm.	
L. Cret.	Albian	Valhall/Rødby Fm.		Unspec. unit	Rødby Fm.		

**Comparison between the Deegan & Scull (1977) and the present subdivision.**

Fig. 6




Fig. 7

CRETACEOUS LITHOSTRATIGRAPHIC NOMENCLATURE  
NORWEGIAN NORTH SEA




Fig. 8

**CRETACEOUS LITHOSTRATIGRAPHIC NOMENCLATURE**  
NORWEGIAN NORTH SEA

R-0189/4





Fig. 9

## CRETACEOUS LITHOSTRATIGRAPHIC NOMENCLATURE NORWEGIAN NORTH SEA

R-0189/4



R-0189/4

Fig. 10

## CRETACEOUS LITHOSTRATIGRAPHIC NOMENCLATURE NORWEGIAN NORTH SEA




Fig.11

CRETACEOUS LITHOSTRATIGRAPHIC NOMENCLATURE  
NORWEGIAN NORTH SEA

R-0189/4

Graben and correlation is therefore difficult. More well data are needed before formal subdivision can be established.

### REVISED CRETACEOUS LITHOSTRATIGRAPHY OF THE NORWEGIAN NORTH SEA

by P. A. Hole, E. Holtar, D. Isaksen, K. S. Lervik, T. Monsen, S. Nybakken, J. E. Tellefsen, K. Tonstad and B.T.G. Wandås.

#### Cromer Knoll Group (Cromer Knollgruppen)

##### Name:

From the Cromer Knoll buoy in the southern North Sea. Named by Deegan & Scull (1977).

##### Type area:

The type area is in the southern North Sea. Rhys (1974) used UK well 48/22-2 to illustrate a typical section of the group, and Deegan & Scull (1977) used UK wells 29/25-1, 22/1-2A and 3/29-1, and Norwegian well 2/11-1. The emphasis in this paper is on the Norwegian sector and the following Norwegian wells have been used to illustrate local developments of the group: 2/6-2, 2/7-15, 2/11-1, 7/3-1, 17/11-2, 34/10-18, 35/3-4, 35/3-5, 31/6-3, 24/12-2 and 17/4-1; the Danish well DK I-1 has also been used.

##### Thickness:

The thickness of the group varies considerably since the sediments were deposited in response to an active Late Jurassic tectonic phase. In the Viking Graben, the Åsta Graben and locally in the Central Trough the thickness is often more than 600 m, gradually thinning towards the basin margins. The group is 667 m thick in Norwegian well 2/11-1 and 643 m thick in Norwegian well 17/11-1. Seismic data indicate that the group is thickest in the Sogn Graben, where it probably reaches up to 1400 m.

##### Lithology:

The Cromer Knoll Group consists mainly of fine-grained, argillaceous, marine sediments with a varying content of calcareous material. Calcareous claystones, siltstones and marlstones dominate, but subordinate layers of limestone and sandstone occur. The claystones are generally light to dark grey, olive-grey, greenish and brownish, often becoming light grey, light greenish-grey and light olive-grey marlstones. Mica, pyrite and glauconite are common. Generally, marlstones become the more dominant lithology in both the upper and lower parts of the group.

##### Basal stratotype:

The lower boundary is usually well defined and is recognised by a distinct decrease in gamma-ray response and an increase in velocity when passing upward from the generally more organic-rich shales of the underlying Upper Jurassic formations (Figs. 12-14 and 22).

##### Characteristics of the upper boundary:

South of approximately 59° N (Fig. 32a), the upper

boundary is the base of the chalk facies of the Shetland Group, defined by the onset of a decrease in gamma-ray response and an increase in velocity into the overlying carbonates (Figs. 12-15 and 22-28). The uppermost Rødby Formation of the Cromer Knoll Group often appears on logs as a transition between the overlying carbonates of the Shetland Group and the more argillaceous parts of the Cromer Knoll Group (Figs. 12 and 22). Further north, the upper boundary is the base of the siliclastic facies of the Shetland Group (Fig. 32a). This boundary is normally also shown by a decrease in gamma-ray response and an increase in velocity when passing into the overlying, generally more calcareous, Svarte Formation of the Shetland Group (Figs. 33 and 36). However, the opposite log response at the boundary is also observed (Fig. 37).

##### Distribution:

The group is widely distributed in the Norwegian sector of the North Sea. It is absent from the highest parts of the Mandal High, Jæren High, Utsira High and Lomre Terrace in the Troll area and locally from the Tampen Spur. (Figs. 7-11).

##### Age:

Ryazanian (usually late) to Albian/Early Cenomanian.

##### Depositional environment:

Open marine, with generally low energy.

##### Subdivision:

Six formations are defined within the group in the Norwegian sector (Figs. 7-11): the Åsgard (new), Tuxen (Jensen et al. 1986), Mime (new), Sola (Hamar & Hesjedal 1983; Jensen et al. 1986), Rødby and Agat (new) Formations. In addition we recognise a need to introduce the Ran sandstone units (new) (Figs. 7-11). Fig. 4 shows an idealised development of the Cromer Knoll Group in the northern and central North Sea.

##### Remarks:

The group was erected by Rhys (1974) to embrace three marine, arenaceous, argillaceous to marly formations of mainly Early Cretaceous age recognisable onshore and offshore. Deegan & Scull (1977) formally defined the group to include the sediments between the underlying Humber Group and Bream Formation and the overlying Shetland and Chalk Groups. Vollset & Doré (1984) replaced the Humber Group of the northern North Sea by the Viking Group, and the Bream Formation in the Central Trough and the Norwegian-Danish Basin by the Tyne and Boknfjorden Groups, respectively. The tops of the Draupne Formation of the Viking Group, the Mandal Formation of the Tyne Group and the Flekkefjord Formation of the Boknfjorden Group define the base of the Cromer Knoll Group.

The Cromer Knoll Group is partly equivalent to the Rijnland Group of the Dutch sector (NAM & RGD 1980, Crittenden 1982) and the Speeton Clay Formation together with the Red Chalk Formation of the UK sector (Rhys 1974). The subdivision in this paper can be used for the Danish sector (see also Jensen et al. 1986).



CRETACEOUS  
WELL 2/11-1

REFERENCE WELL: ÅSGARD FORMATION,  
SOLA FORMATION, RØDBY FORMATION




Fig. 12

CRETACEOUS  
WELL 17/11-2

REFERENCE WELL:  
ÅSGARD FORMATION  
RAN SANDSTONE UNITS




Fig. 13

**CRETACEOUS  
WELL DK I-1**

**TYPE WELL: SOLA FORMATION  
TUXEN FORMATION  
REFERENCE WELL: ÅSGARD FORMATION**




Fig. 14

**CRETACEOUS  
WELL 2/6-2**

**REFERENCE WELL: TUXEN FORMATION**




Fig. 15

### Åsgard Formation (new) (Åsgardformasjonen)

#### Name:

Named from Norse mythology after the castle of the Norse gods, where Odin ruled.

#### Well type section:

Norwegian well 2/11-1 from 3555 to 3063 m, coordinates N 56°14'16.98", E 03°27'07.05" (Fig. 12). No cores.

#### Well reference sections:

Norwegian well 17/11-2 from 2410 to 1802 m, coordinates N 58°06'54.91", E 03°22'09.81" (Fig. 13). No cores.

Danish well I-1 from 3358 to 2986 m, coordinates N 56°03'10", E 04°14'60" (Fig. 14). No cores.

#### Thickness:

The formation is 492 m thick in the type well and 608 m thick in reference well 17/11-2. In the Central Trough area the thickness varies from a few metres to more than 500 m over short distances, showing the complex pattern of small, restricted Early Cretaceous basins. An even thicker sequence was penetrated in the Norwegian-Danish Basin, and especially in the Åsta Graben, where more than 700 m were encountered in well 17/12-3. The formation is thickest in the Sogn Graben where it is probably more than 1200 m, as indicated by seismic data.

#### Lithology:

The formation is dominated by light to dark grey, olive-grey, greenish and brownish, often calcareous claystones, and passes into light grey, light greenish-grey and light olivegrey marlstones and stringers of limestone. Mica, pyrite and glauconite are common. The claystones may be silty, and siltstones or very fine-grained sandstone layers or laminae are present. Where major sandstone layers occur they are regarded as belonging to the Ran sandstone units defined below. In a few Norwegian wells in the central North Sea (e.g. 1/9-3, 2/3-1, 2/7-2, 2/10-1, 2/11-1, 7/3-1, 7/8-2, 7/12-4, 7/12-5 and 8/1-1) a sequence of calcareous claystone, marlstone and limestone interbeds is recognised as the basal part of the Åsgard Formation (Figs. 12 and 14). This sequence is very difficult to correlate in the Norwegian sector, even over small distances, and is therefore regarded as representing local variations in the lowermost part of the Åsgard Formation. In the Danish sector this sequence is defined as the Leek Member (Jensen et al. 1986).

#### Basal stratotype:

The lower boundary is defined by a marked upward decrease in gamma-ray response and an increase in velocity in areas where the underlying sediments are slightly to non-calcareous, organic-rich claystones and shales, usually belonging to the Mandal, Flekkefjord, Tau or Draupne Formations (Figs. 15 and 22). Where the claystones and shales are less organic rich and more calcareous, the boundary may be more difficult to identify on logs.

#### Characteristics of the upper boundary:

The characteristics of the upper boundary vary with the overlying formations. Where the Tuxen Formation occurs, the boundary is defined by an upward decrease

in the gamma-ray readings and an increase in velocity, reflecting slightly more calcareous claystones, marlstones and limestones compared with the underlying Åsgard Formation (Figs. 14 and 15). Where the Tuxen Formation is missing and the Sola Formation is deposited on the Åsgard Formation, the boundary is defined by an upward increase in gamma-ray readings and a decrease in velocity (Figs. 17, 18 and 23). If both the Tuxen and Sola Formations are missing, the boundary to the overlying Rødby Formation is defined by an upward decrease in gamma-ray readings and an increase in velocity. Locally, the Åsgard Formation is overlain by the Ran sandstone units (Fig. 22) and the Agat Formation (Figs. 19 and 20). This boundary is defined by an upward decrease in gamma-ray readings.

#### Distribution:

The Åsgard Formation is very widespread in the North Sea (Figs. 7-11), as are the partial equivalents in the Danish sector (Valhall Formation, Jensen et al. 1986; Vedsted Formation, Larsen 1966), British sector (Speeton Clay, Rhys 1974) and Dutch sector (Vlieland Shale Member, NAM & RGD 1980). In the Norwegian sector, the formation is absent from the highest parts of the Mandal, Jæren and Utsira Highs, the Lomre Terrace, the Troll area, Tampen Spur and locally over salt pillows and diapirs in the Central Trough and the Norwegian-Danish Basin.

#### Age:

Where the Tuxen Formation occurs, the Åsgard Formation ranges in age from Late Ryazanian to Late Hauterivian. In areas where neither the Tuxen nor Sola Formations are recognised, the Åsgard Formation represents a lateral equivalent and may reach Late Aptian to Early Albian age.

#### Depositional environment:

The formation was deposited in an open marine, low-energy shelf environment with well-oxygenated bottom water.

#### Remarks:

Deegan & Scull (1977) divided the Cromer Knoll Group into the Rødby and Valhall Formations. Several lithostratigraphic units have later been described in the Valhall Formation (Hesjedal & Hamar 1983, Jensen et al. 1986). The remaining claystones and marlstones of the originally defined Valhall Formation constitute the Åsgard Formation.

### Tuxen Formation (Tuxenformasjonen)

#### Name:

Named by Jensen et al. (1986) from a bathymetric feature west of Blåvandshuk, the westernmost point of Jylland.

#### Well type section:

Danish well I-1 from 2986 to 2898 m, coordinates N 56°03'10", E 04°14'60" (Fig. 14). No cores.

#### Well reference sections:

Norwegian well 2/11-1 from 3063 to 2988 m, coordinates N 56°14'16.98", E 03°27'07.05" (Fig. 12). No cores.

Norwegian well 2/6-2 from 3935 to 3864 m, coordinates N 56°30'48.90", E 03°42'39.66" (Fig. 15). No cores.

**Thickness:**

The thickness of the formation varies from 1 m along structural highs to about 100 m in basinal areas. In the reference wells the thicknesses are 75 m (2/11-1) and 71 m (2/6-2). In the type well (I-1) the thickness is 88 m.

**Lithology:**

The formation is dominated by white to greyish-pink, calcareous claystones and marlstones. Along some of the structural highs the marlstones grade into purer limestones. Generally, the formation terminates vertically upwards with a chalk sequence containing subordinate marlstone layers. This chalk is white to pale orange or yellowish-grey, occasionally greenish and reddish. The marlstones are generally light grey to greenish-grey or olive-grey, but may be reddish-brown in some wells.

A 0.3-1 m thick, radioactive, marlstone bed is frequently encountered within the Tuxen Formation in the Danish sector where it is defined as the Munk Marl Bed (Jensen et al. 1986). This characteristic unit has also been recognised in some wells in the central Norwegian sector (e.g. 2/1-2, 2/1-3, 2/1-8, 2/6-2, 2/11-7, 6/3-1, 16/8-1 and 16/10-1), (see also Fig. 15). In the Norwegian sector, the Tuxen Formation above the Munk Marl Bed is often more calcareous than the rest of the sequence.

**Basal stratotype:**

The lower boundary is defined as the base of an upward decrease in gamma-ray readings and an increase in velocity, reflecting the passage from the slightly calcareous claystones of the underlying Åsgard Formation up into the more calcareous claystones and marlstones of the Tuxen Formation (Figs. 12, 14 and 15). The transition is generally gradual in basinal areas. Purer limestones were deposited along some structural highs, causing more distinct log breaks.

**Characteristics of the upper boundary:**

Upwards, the Tuxen Formation is generally in contact with the micaceous claystones and organic-rich shales of the Sola Formation (Figs. 12 and 14). This boundary is marked by an upward increase in gamma-ray readings and a decrease in velocity. Where the Sola Formation is missing, the Tuxen Formation is in contact with the marlstones of the overlying Rødby Formation (Fig. 15). The boundary is usually defined by an upward increase in gamma-ray readings.

**Distribution:**

The Tuxen Formation is widely distributed in the Norwegian and Danish sectors (Jensen et al. 1986). In the Norwegian sector it is developed in the Central Trough, along the Jæren High and in parts of the Norwegian-Danish Basin.

In basinal areas in the Norwegian sector it inter-fingers laterally with claystones and marlstones of the Åsgard Formation (Figs. 4 and 7).

**CRETACEOUS  
WELL 34/10-18**

**TYPE WELL: MIME FORMATION**




Fig. 16

**CRETACEOUS  
WELL 17/4-1**

**REFERENCE WELL: MIME FORMATION**




Fig. 17

**Age:**

Late Hauterivian to Late Barremian (Heilmann-Clausen 1987, Thomsen 1987).

**Depositional environment:**

Deposition was dominated by pelagic marl and chalk oozes, which covered large areas of the North Sea. The bottom waters were mainly well oxygenated (Jensen et al. 1986).

**Mime Formation (new) (Mimeformasjonen)****Name:**

Named after a god from Norse mythology who was considered to be very wise.

**Well type section:**

Norwegian well 34/10-18 from 2351 to 2340 m, coordinates N 61°14'22.48", E 02°03'18.83" (Fig. 16). Cores (lower half of the formation).

**Well reference section:**

Norwegian well 17/4-1 from 2122 to 2080 m, coordinates N 58°35'54.00", E 03°16'05.00" (Fig. 17). No cores.

**Thickness:**

In the type well the Mime Formation is 11 m, and in the reference well it is 42 m thick. Usually, the thickness varies between 5 and 20 m.

**Lithology:**

The formation is dominated by limestones and marls. It often contains impure carbonates that are reworked and mixed with smaller quantities of sand and silt. The formation is sometimes chalky. The matrix is usually very calcareous. Oolites are observed in some wells in the East Shetland Basin. The colour is usually white or light pink, but may vary slightly on account of the sand/silt mixture.

**Basal stratotype:**

In those wells where the Mime Formation is present it defines the lower boundary of the Lower Cretaceous, lying on the Upper Jurassic sediments or older rocks. This boundary is always an unconformity and can most often be seen on the logs as a decrease in gamma-ray readings and an increase in velocity upwards from the underlying Jurassic sediments.

**Characteristics of the upper boundary:**

The upper boundary is usually defined at the bottom of more or less calcareous shales in the Åsgard Formation. This boundary is reflected on the logs as an upward increase in gamma-ray readings and a reduction in velocity (Fig. 17). The upper boundary can also be defined by the overlying shales of the Sola or Rødby Formations. The boundary will normally be reflected on logs as described above (Fig. 16).

**Distribution:**

The formation is found only as narrow zones along structural highs. On the flanks of the Viking Graben it may be seen almost continuously from approximately 58° to 62° N. It is also found as a thin carpet over most of the East Shetland Basin and along the Fladen Ground Spur, the Utsira High-Lomre Terrace, the northwest side of the Sele High, and the Jæren High.

The formation is not encountered in the more central parts of the basins, and it is doubtful if it is present along the boundaries of the Fennoscandian Shield.

**Age:**

The formation is time-transgressive, and is dated to Late Valanginian to Albian. It is oldest in the deeper parts along the basin margins and becomes younger up along the flanks. In most of the East Shetland Basin, along the Utsira, Bergen, Sele and Jæren Highs, and along the flanks of the Viking Graben, it is usually of Barremian/Hauterivian age (Fig. 4).

**Depositional environment:**

Palaontological investigations together with the observation of oolites indicate a transgressive, shallow marine, depositional environment.

**Remarks:**

Hesjedal & Hamar (1983) described the impure, reworked limestones resting directly on the Base Cretaceous unconformity over the structural highs, as the Utvik Formation. This formation is formally defined as the Mime Formation in this paper, since the name suggested was not in accordance with existing recommendations.

**Sola Formation (Solaformasjonen)****Name:**

Informally named by Hesjedal & Hamar (1983) after a village in southwestern Norway. Formally named by Jensen et al. (1986).

**Well type section:**

Danish well I-1 from 2898 to 2859 m. Coordinates N 56°03'10", E 04°14'60" (Fig. 14). No cores.

**Well reference sections:**

Norwegian well 2/11-1 from 2988 to 2910 m, coordinates N 56°14'16.98", E 03°27'07.05" (Fig. 12). No cores.

Norwegian well 24/12-2 from 4043 to 3985 m, coordinates N 59°12'00.75", E 01°52'53.34" (Fig. 18). No cores.

**Thickness:**

The thickness in the type well is 39 m, and in reference well 2/11-1 it is 78 m. It generally varies between 20 m and 200 m. The formation is thick in the Viking Graben and Åsta Graben, and thin in the East Shetland Basin and parts of the Fiskebank Sub-Basin.

**Lithology:**

The Sola Formation consists of shales interbedded with stringers of marlstone and limestone. The carbonate content is lower than that in the underlying Tuxen and Åsgard Formations and the overlying Rødby Formation. The colour is black or dark grey, but olive-grey, brown and red colours occur. The shales are finely laminated and often very pyritic.

**Basal stratotype:**

The lower boundary is usually placed on the Tuxen or Åsgard Formations (Figs. 12, 14, 17 and 18). Generally, the gamma-ray response increases and the velocity decreases from the calcareous and sandy sedi-

## CRETACEOUS

## WELL 24/12-2


REFERENCE WELL: SOLA FORMATION  
(CROMER KNOLL GROUP)

Fig. 18

ments up into the shaly and organic rich Sola Formation. In some wells in the east, on the Horda Platform, the gamma-ray response does not increase when the boundary from the Åsgard Formation up into the Sola Formation is crossed. In such wells, a lower, more stable velocity identifies the Sola Formation.

*Characteristics of the upper boundary:*

The upper boundary is most often placed where the carbonate content starts to increase rapidly into the overlying Rødby Formation (Figs. 12, 14, 18, and 22). In some areas where, the Sola Formation is overlain by the Ran sandstone units (Fig. 23), the boundary is defined by an upward decrease in gamma-ray response and an increase in velocity.

*Distribution:*

The formation is widespread in the North Sea. It is absent or thin on structural highs, salt-induced structu-

res and in parts of the Central Trough and Norwegian-Danish Basin.

*Age:*

The Sola Formation is of Mid Aptian-Early Albian age. A possible Middle (Late) Barremian-Albian age is recorded from Danish wells (Heilmann-Clausen 1986).

*Depositional environment:*

The Sola Formation was deposited in a marine environment with alternating anoxic and oxic bottom conditions. Hesjedal & Hamar (1983) suggested that the formation was deposited during a regressive period, while Rawson & Riley (1982) held the opposite view.

**Rødby Formation (Rødbyformasjonen)***Name:*

Named by Larsen (1966) after a town on the island of Lolland in southern Denmark.

*Well type section:*

The Danish well Rødby-1 drilled on the island of Lolland (Larsen 1966).

*Well reference sections:*

Norwegian well 2/11-1 from 2910 to 2887 m, coordinates N 56°14'16.98", E 03°27'07.05" (Fig. 12). No cores.

Norwegian well 2/7-15 from 3419 to 3401 m, coordinates N 56°23'46.82", E 03°18'54.63" (Fig. 22). No cores.

*Thickness:*

In the well type section the thickness is 23 m. The thicknesses in the reference sections are 23 m in well 2/11-1 and 18 m in well 2/7-15. The formation generally ranges in thickness between 15 and 30 m. In the Viking Graben it may reach thicknesses of more than 200 m (Deegan & Scull 1977).

*Lithology:*

Mainly red-brown marlstones, but green and grey colours may occur. Glauconite and pyrite may be present. Sandstones and siltstones are known to be present locally.

*Basal stratotype:*

The lower boundary is placed on the Sola and Åsgard Formations and the Ran sandstone units, and represents an upward decrease in gamma-ray response and usually an increase in velocity into the Rødby Formation (Figs. 12, 22 and 23).

*Characteristics of the upper boundary:*

The upper boundary can be seen as an upward decrease in gamma-ray response and a increase in velocity when going into the more calcareous sediments of the Svarte Formation (Figs. 33-36). On the Horda Platform the Rødby Formation is more calcareous and has a lower gamma-ray response and a higher velocity than the overlying unspecified unit of the Shetland Group (Fig. 37). Where the chalk facies of the Shetland Group is present, the upper boundary is characterised by a distinct upward drop in gamma-ray readings and a marked increase in velocity (Figs. 12, 15, 22-28).

CRETACEOUS

WELL 35/3-4

TYPE WELL: AGAT FORMATION




Fig. 19

CRETACEOUS

WELL 35/3-5

REFERENCE WELL: AGAT FORMATION




Fig. 20

**Distribution:**

The formation is widespread in the North Sea, but are missing locally on the Utsira, Jæren and Mandal Highs, Tampen Spur and Horda Platform on the Norwegian sector.

**Age:**

The formation is of Albian age (locally Early Cenomanian in the Danish sector (Jensen et al. 1986)).

**Depositional environment:**

The Rødby Formation was deposited as generally reddish sediments in an open marine, oxygenated environment with limited supply of clastics.

**Agat Formation (new) (Agatformasjonen)****Name:**

Named after the gas-condensate Agat Field in Norwegian block 35/3.

**Well type section:**

Norwegian well 35/3-4 from 3589 to 3345 m, coordinates N 61°51'54.54", E 03°52'26.99" (Fig. 19). 95 m of cores, mainly from the lower half of the formation.

**Well reference section:**

Norwegian well 35/3-5 from 3620 to 3219 m, coordinates N 61°47'46.71", E 03°54'44.01" (Fig. 20). 65 m of cores from the upper part of the formation.

**Thickness:**

In the type well the gross thickness of the formation is 244 m, and in the reference well 401 m. The gross thickness varies in that range in the 35/3 wells.

**Lithology:**

In the type well the formation consists of white to light grey, fine- to medium-grained, moderately to well-sorted sandstones alternating with grey claystones. The sandstones are usually micaceous and glauconitic and sometimes contain small amounts of pyrite. The sandstones in the type well are carbonate- and silica-cemented in zones. In the reference well, the upper part of the formation consists of medium- and coarse-grained to pebbly sandstones and conglomerates alternating with dark grey claystones. The conglomerates are both matrix- and grain-supported. The claystones are often found as 0.5-5 m thick layers between the sandstones. They are dark grey, usually calcareous and contain varying amounts of siltstone. They may occasionally pass into light grey, micaceous, calcareous and glauconitic siltstones.

**Basal stratotype:**

The lower boundary is defined where sandstones become the dominant lithology and is placed at the base of the first marked coarsening-upwards sandstone unit or distinct sand body. On logs it shows as an upward reduction in gamma-ray response (Figs. 19 and 20) and most often an increase in velocity (Fig. 20).

**Characteristics of the upper boundary:**

The upper boundary is placed at the top of the upper sandstone layer. This boundary is especially distinct on the gamma-ray log since the overlying sediments are dominated by calcareous shales with a low sandstone content. The overlying sediments are represented either

by the Rødby Formation (well 35/3-1 and 2), or by the Svarte Formation (wells 35/3-4 and 5), (Figs. 19 and 20).

**Distribution:**

The formation is encountered in the area around the Måløy Fault Blocks in Norwegian blocks 35/3-36/1 (Fig. 21) and is expected to be present along the western boundary of the Fennoscandian Shield. It is assumed to pass into shales towards the west (Fig. 21).

**Age:**

Aptian-Albian (possibly Early Cenomanian).

**Depositional environment:**

Marine environment influenced by gravity flows of sediment.

**Remarks:**

Hesjedal & Hamar (1983) named the sand lobes and distal fan delta sands in the Agat Field, and a possible elongation (or similar development) along the Øygarden Fault Zone, as the Florø Formation. This formation is formally defined as the Agat Formation in this paper, since the name suggested does not conform with existing recommendations.

**Ran sandstone units (new) (Ransandsteinsenhetene)****Name:**

Ran was the wife of the sea god Gir in Norse mythology. She liked to drag sailors down to the depths with her net.




Fig. 21

**Well type section:**

None.



**CRETACEOUS**  
**WELL 2/7-15**  
REFERENCE WELL:  
**RAN SANDSTONE UNITS**  
(CROMER KNOLL GROUP)




Fig. 22

**Well reference sections:**

Norwegian well 2/7-15 from 3498 to 3450 m, coordinates N 56°23'46.82", E 03°18'54.63" (Fig. 22). 16 m of cores in the lowermost part of the formation.

Norwegian well 7/3-1 from 2412 to 2396 m, coordinates N 57°50'35.25", E 02°44'55.61" (Fig. 23). No cores.

Norwegian well 17/11-2 from 1802 to 1767 m, coordinates N 58°06'54.91", E 03°22'09.81" (Fig. 13). No cores.

**Thickness:**

The gross sandstone thicknesses vary from a few metres up to approximately 100 m. The gross thicknesses in the reference wells are 48 m (2/7-15), 16 m (7/3-1) and 35 m (17/11-2). Up to 130 m (gross) of Aptian-Albian sandstone sequences are penetrated in block 16/27 in the UK sector (see Distribution).

**CRETACEOUS**  
**WELL 7/3-1**  
REFERENCE WELL:  
**RAN SANDSTONE UNITS**  
(CROMER KNOLL GROUP)




Fig. 23

**Lithology:**

The colour of the sandstones ranges from white to light grey, green and brown to reddish-brown. The sandstones are generally argillaceous, sometimes calcareous and glauconitic, and usually do not represent potential reservoir rocks in these wells.

**Basal stratotype:**

The various sandstone units may appear in contact with the Åsgard, Tuxen, Sola and Rødby Formations (Figs. 4,7,8 and 9). Their lower boundaries are generally defined as the base of an upward decrease in the gamma-ray response when passing into the sandstone units (Figs. 22 and 23). The gamma-ray readings in the calcareous marlstones and chalks of the Tuxen Formation, especially its upper part, and the Mime Formation may be similar to those in the sandstones. The velocity curve is often less suitable for defining the lower boundary.


**CRETACEOUS**

**WELL 1/3-1**

TYPE WELL: HIDRA FORMATION  
HOD FORMATION  
TOR FORMATION

REFERENCE WELL: EKOFISK FORMATION  
BLODØKS FORMATION

(SHETLAND GROUP)



OVERLAPS WITH SECTION BELOW




Fig. 24

*Characteristics of the upper boundary:*

The upper boundary can usually be identified as an upward increase in the gamma-ray readings (Fig. 22) and generally by a slight decrease in the sonic velocity.

*Distribution:*

The Ran sandstone units are encountered in only a few wells in the Norwegian sector (Fig. 21 and Remarks).

*Age:*

Ryazanian-Albian.

*Depositional environment:*

The sandstones that have been penetrated are described as shallow (Norwegian sector) and deep water (UK sector) submarine fans.

*Remarks:*

Hesjedal & Hamar (1983) recognised several scattered sandstone sequences which they described as the Kopervik and Klepp Formations in the Central Trough and Norwegian-Danish Basin, and the Florø Formation in the Agat Field in blocks 35/3 and 36/1. The Kopervik and Klepp Formations are here described as the Ran sandstone units. Since they consisted of several isolated sandstone bodies they should not have been given formation status, and the names did not conform with the existing recommendations. The Florø Formation is formally defined as the Agat Formation in this paper (see also General lithostratigraphic notes for Cretaceous).

In the UK sector (the Andrew Field), just south of the Andrew Ridge and Fladen Ground Spur, Aptian-Albian sandstone sequences (the Bosun Member) are encountered in many wells, among others UK wells 16/27-1 and 16/27a-2 (100-130 m gross), 16/28-1 (50 m gross) and 16/28-6 (90 m gross). The palaeogeographical position of these sandstones, i.e. basinal areas close to the subaerially exposed major structural highs mentioned above, may be quite similar to the palaeogeographical situation along the western margin of the Måløy Fault Blocks. Here, up to 400 m (gross) thick sandstone sequences of Aptian-Early Cenomanian age were deposited in Norwegian blocks 35/3 and 36/1, and are defined as the Agat Formation in this paper.

The Devil's Hole Formation (UK well 29/25-1) and the "Unnamed Formation" (UK well 14/20-5) in the UK sector are comparable to the Ran sandstone units.

**Shetland Group (Shetlandsgruppen)***Name:*

Named from the Shetland Islands off the north coast of Scotland (Deegan & Scull 1977). The group has now been expanded to include the formations of the former Chalk Group.

*Type area:*

The group is typically developed in the central and Northern North Sea. A chalk facies is developed in the central North Sea and a siliclastic facies in the Northern North Sea (Fig. 32 a). A typical section of the chalk facies in the central area is represented by Norwegian well 1/3-1 (Fig. 24), while Norwegian well 25/1-1 (Fig. 33) provides a typical section of the siliclastic facies in the northern area. UK well 22/1-2A illustrates a section in the transition zone between the two facies (Figs. 5 and 25).

*Thickness:*

In well 1/3-1 the group is 1183 m thick, and in well 25/1-1 it measures 1284 m. Seismic interpretation and well data indicate that the thickness of the group ranges between 1000 and 2000 m in the graben areas. The group shows considerable thinning towards and in the platform areas.

*Lithology:*

The group consists of the chalk facies of chalky limestones, limestones, marls, and calcareous shales and mudstones. Chert (flint) occurs throughout the facies. The siliclastic facies consists of mudstones and shales, partly interbedded with limestones. Minor amounts of sandstones are present in the lower part in the Agat Field area (block 35/3). The shales and sandstones are slightly to very calcareous. In the Maastrichtian sequence the quantity of limestones are generally higher on the Horda Platform than in the Viking Graben.

*Basal stratotype:*

Typically the lower boundary is the contact to the calcareous mudstones or marlstones of the Cromer Knoll Group. On structural highs like the Horda Platform, Tampen Spur, Sørvestlandet and Mandal Highs the lower part of the group is occasionally absent and the remainder rests unconformably on the Cromer Knoll Group, Jurassic or older rocks.

*Characteristics of the upper boundary:*

The group is overlain by Paleocene mudstones, marls or sandstones of the Rogaland Group.

*Distribution:*

The group is present throughout the Norwegian North Sea, being absent only locally on highs (e.g. 16/5-1, 31/2-9) and a few salt diapirs (e.g. 2/7-12). A transition between the chalk and siliclastic facies of the group occurs relatively abruptly in the Norwegian sector along the Utsira High (Figs. 32a and b) and more gradually in the graben areas.

*Age:*

The group ranges in age from Cenomanian to Danian. The siliclastic facies is restricted in age to the Late Cretaceous.

*Depositional environment:*

The Late Cretaceous sequence in the North Sea was deposited in an open marine environment during a general rise in sea level (Hancock & Kauffman 1979). The chalk facies formations were deposited as coccolith debris and other carbonate grains and sequences often show a cyclic pelagic sedimentation pattern termed periodite (d'Heur 1986). In the Central Trough, extensive subsidence resulted in the chalk facies being dominated by allochthonous, redeposited chalks which were transported downslope as major slides, slumps, debris flows, and proximal and distal turbidites. The siliclastic facies is less well studied. The influx of siliclastic mud was higher and the carbonate production probably lower than in the area with chalk facies.

*Subdivision:*

The Shetland Group is represented by four chalk facies formations: the Hidra, Hod, Tor and Ekofisk

Formations (all erected by Deegan & Scull 1977) and six siliciclastic facies formations: the Svarte (new), Blodøks (new), Tryggvason (new), Kyrre (new), Jorsalfare (new) and Hardråde (new) Formations.

**Remarks:**

The Herring and Flounder Formations in the UK sector (Deegan & Scull 1977) are regarded as equivalents of the lowermost part of the Hod Formation and of the Kyrre Formation, respectively (Fig. 5 and 6).

**Hidra Formation (Hidraformasjonen)**

**Name:**

Named by Deegan & Scull (1977) after the Hidra High in Norwegian blocks 1/3 and 2/1. The name Hidra is after the island of Hidra on the southern coast of Norway.

**Well type section:**

Norwegian well 1/3-1 from 4441 to 4371 m, coordinates N 56°51'21.00", E 02°51'05.00" (Fig. 24). No cores.

**Well reference sections:**

UK well 22/1-2A from 3783 to 3738 m, coordinates N 57°56'12.20", E 01°02'55.80" (Fig. 25). No cores.

UK well 29/25-1 from 2258.5 to 2228 m, coordinates N 56°18'10.00", E 01°51'48.80" (Fig. 26). No cores.

Danish well BO-1 from 2275.5 to 2220 m, coordinates N 55°48'8.22", E 04°34'18.66" (Fig. 27). Cored through the upper 35 m.

**Thickness:**

The formation is 70 m thick in the type well, 45 m in 22/1-2A, 30.5 m in 29/25-1 and 55.5 m in BO-1. Seismic interpretation suggests that the formation reaches a maximum thickness of about 150 m in the northwestern part of the Central Trough in the Norwegian sector.

**Lithology:**

In the type well the formation consists of white to light grey, hard chalks with thin interbeds of grey to black shale in the lower part of the formation. Locally the formation is more marly with interbedded marly chalk and marl. The chalks are occasionally softer with abundant glauconite and pyrite. The colour may be white, grey, green, brown or pink. At the base of the formation in UK well 22/1-2A, hard, black, carbonaceous and argillaceous limestones are present. Traces of pink waxy tuff occur in places. The formation is generally highly bioturbated.

**Basal stratotype:**

The formation usually shows a gamma-ray response that has constant low values and high velocities. These contrast sharply at the lower boundary with the higher gamma-ray response and lower velocity of the Åsgård and Sola Formations. The lower boundary is more gradational when the carbonate rich facies of the Rødby Formation is present beneath the Hidra Formation.

**Characteristics of the upper boundary:**

The upper boundary is defined by the stratotype of the Blodøks Formation. The boundary is characterised by a change from the chalk lithology to mainly mudstone.

This is seen as an abrupt change to higher gamma-ray response and a decrease in velocity in the Blodøks Formation. The boundary shows as a glauconitised hardground in the core from Danish well BO-1.

**Distribution:**

The formation is found in the central and southern North Sea. In the Norwegian sector, it is missing above highs such as the Sørvestlandet, Mandal, Jæren, Utsira and Sele Highs, the Grensen Ridge, as well as many of the salt diapirs.

**Age:**

Cenomanian.

**Depositional environment:**

Open marine with a perioditic or turbiditic origin for the sediments.

**Hod Formation (Hodformasjonen)**

**Name:**

Named by Deegan & Scull (1977) from the Hod Field in Norwegian block 2/11. The name Hod derives from one of the twelve principal gods in Norse mythology. Hod was a son of Odin.

**Well type section:**

Norwegian well 1/3-1 from 4343 to 3828 m, coordinates N 56°51'21.00", E 01°51'05.00" (Fig. 24). No cores.

**Well reference sections:**

UK well 29/25-1 from 2225 to 2012 m, coordinates N 56°18'10.00", E 01°51'48.80" (Fig. 26). No cores.

Norwegian well 2/8-8 from 2601 to 2494 m, coordinates N 56°16'50.28", E 03°24'15.93" (Fig. 28). 36 m of cores discontinuously through the upper 78 m and lowermost 6 m of the formation.

**Thickness:**

The formation is 515 m thick in the type well, 213 m in UK well 29/25-1 and 107 m in Norwegian well 2/8-8. In the Norwegian sector, seismic interpretation indicates that the formation may reach a thickness of more than 700 m in the northwestern part of the Central Trough.

**Lithology:**

In the type well the formation consists of hard, white to light grey, crypto- to microcrystalline limestones which may become argillaceous or chalky in places. White, light grey to light brown, soft to hard chalk facies may dominate the formation or alternate with limestones. The limestones may be pink or pale orange. Thin, silty, white, light grey to green or brown, and soft, grey to black, calcareous clay/shale laminae are occasionally present. Pyrite and glauconite may occur throughout the formation and the latter may be common in the lower part.

**Basal stratotype:**

The lower boundary is usually marked by a distinct log break to a lower gamma-ray response and higher velocity from the Blodøks Formation to the Hod Formation (Fig. 24). The boundary may be less distinct when the Blodøks Formation is more calcareous (Fig. 31).

CRETACEOUS  
WELL UK 22/1-2A

REFERENCE WELL: HIDRA FORMATION  
TOR FORMATION  
EKOFISK FORMATION  
(SHETLAND GROUP)




Fig. 25

CRETACEOUS  
WELL UK 29/25-1

REFERENCE WELL: HIDRA FORMATION  
HOD FORMATION  
TOR FORMATION  
(SHETLAND GROUP)




Fig. 26

## CRETACEOUS WELL DK BO-1

REFERENCE WELL: HIDRA FORMATION  
BLODØKS FORMATION  
(SHETLAND GROUP)




Fig. 27

### Characteristics of the upper boundary:

The upper boundary towards the Tor Formation is generally marked by a change in gamma-ray readings to a more constant and slightly lower level, and also by higher velocity, (Figs. 24 and 31). The upper boundary may represent an unconformity in the Ekofisk area (e.g. Norwegian well 2/8-8, Fig. 28).

### Distribution:

The formation is widely distributed in central and eastern parts of the central North Sea, passing laterally into sediments of the Herring and Flounder Formations to the west and the Tryggvason and Kyrre Formations to the northwest.

**Age:**  
Turonian to Campanian.

### Depositional environment:

Open marine with deposition of cyclic pelagic carbonates (periodites) and distal turbidites (Skovbro 1983 and d'Heur 1986).

### Remarks:

An informal, tripartite subdivision of the Hod Formation into lower, middle and upper members is often possible in the southern part of the Central Trough in the Norwegian sector. The subdivision is based on the frequent presence of a higher clay content in the middle of the Hod Formation (Figs. 24, 28 and 29).

### LOWER MEMBER OF THE HOD FORMATION:

This unit constitutes the largest part of the Hod Formation and is a sequence of bioturbated laminated chalks with a low clay content. It occurs in Norwegian wells 1/3-1 from 4343 to 4066 m, 2/8-8 from 2601 to 2538 m and 1/9-1 from 3648 to 3353 m.

### MIDDLE MEMBER OF THE HOD FORMATION:

This is a sequence consisting mainly of periodites, which generally have a greyish colour reflecting a marked increase in terrigenous clay. It is shown on well logs as an increase in gamma-ray readings. It occurs in Norwegian wells 1/3-1 from 4066 to 4009 m, 2/8-8 from 2538 to 2518 m and 1/9-1 from 3353 to 3344 m.

### UPPER MEMBER OF THE HOD FORMATION:

This unit constitutes another sequence dominated by periodites with minor allochthonous intercalations, but with a return to a low clay content. It occurs in Norwegian wells 1/3-1 from 4009 to 3828 m, 2/8-8 from 2518 to 2494 m and 1/9-1 from 3344 to 3312 m.

The Herring Formation of Deegan & Scull (1977) includes a similar lithology and was deposited at the same time as the Hod Formation. It is regarded here as the lower part of the Hod Formation. The Hod Formation is also equivalent in age to the Tryggvason and Kyrre Formations (Fig. 6).

### Tor Formation (Torformasjonen)

#### Name:

Named by Deegan & Scull (1977) from the Tor Field in Norwegian blocks 2/4 and 2/5. Tor was a son of Odin, and one of the principal gods of Norse mythology.

#### Well type section:

Norwegian well 1/3-1 from 3828 to 3354 m, coordinates N 56°51'21.00", E 02°51'05.00" (Fig. 24). No cores.

#### Well reference sections:

UK well 22/1-2A from 3245 to 2982.5 m, coordinates N 57°56'12.20", E 01°12'55.80" (Fig. 25). No cores.

UK well 29/25-1 from 2212 to 1869 m, coordinates N 56°18'10.00", E 01°51'48.80" (Fig. 26). No cores.

Norwegian well 1/9-1 from 3312 to 3104 m, coordinates N 56°24'05.07", E 02°54'06.49" (Fig. 29). Cored through the formation.

**Thickness:**

The formation is 474 m thick in the type well, 262.5 m in 22/1-2A, 143 m in 29/25-1 and 208 m in 1/9-1. In the Norwegian sector, seismic interpretation indicates that the thickness of the formation may exceed 600 m in the northwestern part of the Central Trough.

**Lithology:**

In the type well the formation consists of white to light grey, tan to pink, hard, chalky limestones. The formation is generally homogenous, or consists of alternating white, grey or beige, moderately hard to very hard, rarely soft, mudstones or wackestones, rarely packstones, chalks, chalky limestones or limestones. Occasional fine layers of soft grey-green or brown marl occur and also rare stringers of grey to green calcareous shales.

**Basal stratotype:**

The lower boundary is generally marked by an upward change to a more constant lower level of gamma-ray response, and also by higher velocity (Fig. 24).

**Characteristics of the upper boundary:**

The upper boundary is marked by the end of the more constant low gamma-ray response with a return to a higher and more irregular gamma ray and a lower velocity in the overlying Ekofisk Formation (Fig. 24). The upper boundary represents an unconformity with a submarine hardground along the Lindesnes Ridge, and a change of deposition to clay-rich chalks or minor shales (Fig. 28).

**Distribution:**

The formation is present throughout the central North Sea (Fig. 32b). In the Norwegian sector it is very thin or absent on the Lindesnes Ridge and the Utsira High.

**Age:**

Late Campanian to Maastrichtian.

**Depositional environment:**

Open marine with deposition of calcareous debris flows, turbidites and autochthonous periodites.

**Remarks:**

In general the formation shows an upward increase in the intensity and thickness of allochthonous beds. 16 major correlative allochthonous units are recognised within the Central Trough area (Hatton 1986). The sequence of single or stacked mass flows with autochthonous periodite facies enables three separate members to be distinguished (d'Heur 1984, Hatton 1986, Figs. 24, 29, 30 and 31). The semi-regional distribution pattern of the allochthonous units shows that this subdivision is not valid throughout the area of the chalk facies.

**Lower member of the Tor Formation:**

The sequence is dominated by autochthonous periodite deposits interrupted by single or stacked minor debris flows. The sequence is present in Norwegian wells 2/5-1 from 3475 to 3235 m, 1/3-1 from 3828 to 3447 m and 2/4-5 from TD to 3283 m.

**Middle member of the Tor Formation:**

This unit shows a marked increase in slumps, slides, and stacked and single debris flows relative to intervening minor autochthonous beds. The unit is present in Norwegian wells 2/5-1 from 3235 to 3192 m, 1/3-1 from 3447 to 3411 m, 2/4-5 from 3283 to 3237 m and 1/9-1 from 3168 to 3137 m.

**Upper member of the Tor Formation:**

This unit generally consists of high porosity, homogenous chalks which represent stacked sequences of, often slumped, debris flows. The unit is present in Norwegian wells 2/5-1 from 3192 to 3132 m, 2/4-5 from 3237 to 3164 m and 1/3-1 from 3411 to 3354 m.

**Ekofisk Formation (Ekofiskformasjonen)**

**Name:**

Named after the Ekofisk Field in Norwegian block 2/4 (Deegan & Scull 1977).

**Well type section:**

Norwegian well 2/4-5 from 3164 to 3037 m, coordinates N 56°34'29.77", E 03°12'13.03" (Fig. 30). No cores.

**Well reference sections:**

Norwegian well 1/3-1 from 3354 to 3258 m, coordinates N 56°51'21.00", E 02°51'05.00" (Fig. 24). No cores.

UK well 22/1-2A from 2982.5 to 2935 m, coordinates N 57°56'12.20", E 01°02'55.80" (Fig. 25). No cores.

Norwegian well 2/5-1 from 3132 to 3041 m, coordinates N 56°38'19.95", E 03°21'07.94" (Fig. 31). Cored through the upper 78 m.

**Thickness:**

The formation is 127 m thick in the type well, 96 m in 1/3-1, 47.5 m in 22/1-2A and 91 m in 2/5-1. In the Norwegian sector, seismic interpretation indicates that a thickness of more than 150 m is found in the northwestern part of the Central Trough.

**Lithology:**

In the type well, the formation comprises white, tan or beige, hard, dense, sometimes finely crystalline limestones, although softer chalky textures are also present. The formation usually consists of white to light grey, beige to brownish, mudstones or wackestones with occasional packstones/grainstones and pisolitic horizons, often alternating with argillaceous chalks, chalky limestones or limestones. Thin beds of grey, calcareous, often pyritic shales or clays are most common in the lower part while brownish-grey cherts occur rarely to abundantly throughout the formation.

**Basal stratotype:**

The lower boundary is marked by a change in gamma-ray readings from a constant low level in the Tor Formation to a slightly lower level. The velocity may or may not show a corresponding increase. The lower boundary separates the Cretaceous and Tertiary chalks and may represent an unconformity (e.g. Norwegian well 1/9-1, Fig. 29).

CRETACEOUS  
WELL 2/8-8

REFERENCE WELL: HOD FORMATION  
(SHETLAND GROUP)




Fig. 28

CRETACEOUS  
WELL 1/9-1

REFERENCE WELL: TOR FORMATION  
(SHETLAND GROUP)




Fig. 29



CRETACEOUS

WELL 2/4-5

TYPE WELL: EKOFISK FORMATION

(SHETLAND GROUP)




Fig. 30

CRETACEOUS

WELL 2/5-1

REFERENCE WELL: EKOFISK FORMATION

(SHETLAND GROUP)




Fig. 31

**Characteristics of the upper boundary:**

The upper boundary is defined where the gamma-ray response increases and the velocity decreases towards the marly beds of the Våle Formation. Where the marl is not present the change is more abrupt (e.g. Norwegian well 2/8-8, Fig. 28).

**Distribution:**

The formation is widespread in the southern and central North Sea. In the Norwegian sector, it is missing from parts of the Sørvestlandet High and the Lindesnes Ridge.

**Age:**

Danian.

**Depositional environment:**

Open marine with deposition of calcareous debris flows, turbidites and autochthonous periodites (Skovbro 1983, d'Heur 1986, Hatton 1986).

**Remarks:**

Two zones of the formation are readily correlatable within the Central Trough area (Hatton 1986, Figs. 24, 29, 30 and 31).

**LOWER MEMBER OF THE EKOFISK FORMATION:**

The lowermost part consists of a low porosity to tight zone with a higher terrigenous clay content, and is informally termed the Ekofisk tight zone. The larger part consists of the informal Ekofisk reworked zone with mainly reworked Maastrichtian chalks (Tor

Formation) deposited as various mass flows and periodite-facies chalks. This lower member is present in Norwegian wells 1/3-1 from 3354 to 3307 m, 1/9-1 from 3104 to 3072 m, 2/4-5 from 3164 to 3106 m and 2/5-1 from 3132 to 3099 m.

**UPPER MEMBER OF THE EKOFISK FORMATION:**

This zone is composed of mainly homogenous chalks with a low clay content, debris flows of reworked Danian chalks and minor turbidites. A lower tight to low porosity zone, informally termed the Tommeliten tight zone, is present in parts of the Central Trough. The zone is found in Norwegian wells 1/3-1 from 3311 to 3258 m, 1/9-1 from 3072 to 3036 m, 2/4-5 from 3102 to 3037 and 2/5-1 from 3094 to 3041 m.

**Svarte Formation (new) (Svarteforrasjonen)****Name:**

Named after Halvdan Svarte, King of Ringerike, Norway about A.D. 850.

**Well type section:**

Norwegian well 25/1-1 from 3995 to 3807 m, coordinates N 59°53'17.40", E 02°04'42.70" (Fig. 33). No cores.

**Well reference sections:**

Norwegian well 35/3-2 from 3447 to 3207 m, coordinates N 61°51'05.98", E 03°46'28.22" (Fig. 34). No cores.

Norwegian well 24/9-1 from 3992 to 3804 m, coordinates N 59°16'09.48", E 01°47'31.18" (Fig. 35). No cores.

Approximate boundary between the Shetland Group. Siliclastic facies and chalk facies.




Fig. 32 a

Approximate boundary between the Jorsalfare, Hardråde and Tor Formations.




Fig. 32 b

**Thickness:**

In the Viking Graben, the formation is 188 m thick in the type well 25/1-1, 240 m in well 35/3-2 and 188 m in well 24/9-1.

**Lithology:**

The formation generally consists of mudstones interbedded with limestones. Sandstones occur in the Agat area. The content of limestones relative to mudstones is generally lower in the northern than in the southern part of the Viking Graben. The mudstones are medium to light grey, often calcareous, occasionally micaceous, glauconitic and pyritic. The limestones are mainly white to medium grey, argillaceous or sandy. The sandstones are clear to light grey and often cemented by calcite.

**Basal stratotype:**

The lower boundary shows a general upward decrease in gamma-ray intensity and an increase in velocity from the Cromer Knoll Group into the Svarte Formation (Fig. 34). This is due to a higher content of carbonate in the Svarte Formation.

**Characteristics of the upper boundary:**

The upper boundary is generally easily located, and is characterised by an increase in gamma-ray intensity and a distinct decrease in velocity from the Svarte Formation up into the Blodøks Formation (Fig. 33). This is caused by a lower carbonate content in the Blodøks Formation.

**Distribution:**

The formation is present in the Viking Graben and north of the Tampen Spur towards the Marulk Basin. It is, however, lacking on structural highs such as the Lomre Terrace (e.g. Norwegian wells 35/8-1 and 35/8-2), (Nybakken and Bäckström, in press).

**Age:**

Cenomanian.

**Depositional environment:**

Open marine.

**Remarks:**

The Svarte Formation is time-equivalent with the Hydra Formation in the central North Sea and with the informal "formation A" of Deegan & Scull (1977) (Fig. 6).

**Blodøks Formation (new) (Blodøksformasjonen)****Name:**

Named after Eirik Haraldson Blodøks, a Norwegian king who reigned in Norway (A.D. 930-934) and in Northumberland (A.D. -954).

**Well type section:**

Norwegian well 25/1-1 from 3807 to 3790 m, coordinates N 59°53'17.40", E 02°04'42.70" (Fig. 33). No cores.

**Well reference sections:**

Norwegian well 35/3-2 from 3207 to 3190 m, coordinates N 61°51'05.98", N 03°46'28.22" (Fig. 34). No cores.

Norwegian well 1/3-1 from 4371 to 4343 m, coordinates N 56°51'21.00", E 02°51'05.00" (Fig. 24). No cores.

Danish well BO-1 from 2220 to 2213 m, coordinates N 55°48'02.22", E 04°34'18.66" (Fig. 27). Cored throughout.

**Thickness:**

The formation is 17 m thick in the type well (25/1-1), 17 m in well 35/3-2, 28 m in well 1/3-1 and 7 m in well BO-1. It rarely exceeds 20 m in thickness.

**Lithology:**

The formation consists of red, green, grey and black shales and mudstones which are non-calcareous to moderately calcareous. In the central North Sea the formation may show a varied influx of marls, limestones and chalky limestones.

**Basal stratotype:**

The lower boundary is generally characterised by a distinct log break with an upward increase in gamma-ray intensity and a distinct decrease in velocity from the Svarte Formation (Fig. 33) or Hydra Formation (Fig. 24) into the Blodøks Formation. This is due to the lower content of carbonate in the Blodøks Formation.

**Characteristics of the upper boundary:**

The upper boundary shows a decrease in gamma-ray intensity and an increase in velocity from the Blodøks Formation upwards into the more calcareous Tryggvason Formation (Fig. 34) or the chalky Hod Formation (Fig. 24).

**Distribution:**

The formation is present throughout the North Sea, lacking only on local highs such as the Sørvestlandet High, the Utsira, Mandal, Jæren and Sele Highs and the Grensen Ridje as well as above many salt diapirs.

**Age:**

Latest Cenomanian to early Turonian.

**Depositional environment:**

The formation was deposited during a period characterised by anoxic bottom conditions (e.g. Hart & Leary 1989). Presence of carbonates may indicate periods of more oxic conditions or supply of allochthonous limestones and chalks (e.g. Norwegian wells 1/3-1 and 2/5-1).

**Remarks:**

The Blodøks Formation is equivalent to the former Plenus Marl Formation and the informal "formation B" of Deegan & Scull (1977) (Fig. 6). A black shale of Early Turonian age is also widespread outside the North Sea, e.g. the Yorkshire Black Band in England (Jeffries 1963) and similar facies on Helgoland and in northwestern Germany (Schmid & Spaeth 1980).

**Tryggvason Formation (new)  
(Tryggvasonformasjonen)****Name:**


Named after Olav Tryggvason, a Norwegian king (A.D. 995-1000).

CRETACEOUS

WELL 25/1-1

TYPE WELL: SVARTE FORMATION  
 BLODØKS FORMATION  
 TRYGGVASON FORMATION  
 KYRRE FORMATION  
 JORSALFARE FORMATION

(SHETLAND GROUP)



OVERLAPS WITH SECTION BELOW





Fig. 33

CRETACEOUS  
WELL 35/3-2

REFERENCE WELL: SVARTE FORMATION  
BLODØKS FORMATION  
TRYGGVASON FORMATION  
KYRRE FORMATION  
JORSALFARE FORMATION

(SHETLAND GROUP)



OVERLAPS WITH SECTION BELOW





Fig. 34

CRETACEOUS  
WELL 24/9-1

REFERENCE WELL: SVARTE FORMATION  
TRYGGVASON FORMATION  
KYRRE FORMATION  
JORSALFARE FORMATION



OVERLAPS WITH SECTION BELOW




Fig. 35

*Well type section:*

Norwegian well 25/1-1 from 3790 to 3582 m, coordinates N 59°53'17.40", E 02°04'42.70" (Fig. 33). No cores.

*Well reference sections:*

Norwegian well 35/3-2 from 3190 to 2864 m, coordinates N 61°51'05.98", E 03°46'28.22" (Fig. 34). No cores.

Norwegian well 24/9-1 from 3783 to 3638 m, coordinates N 59°16'09.48", E 01°47'31.18" (Fig. 35). No cores.

Norwegian well 30/11-3 from 3207 to 3162 m, coordinates N 60°02'38.59", E 02°32'15.47" (Fig. 36). No cores.

*Thickness:*

In the Viking Graben, the formation is 208 m thick in the type well (25/1-1), 326 m in well 35/3-2 and 145 m in well 24/9-1. It is 45 m thick in well 30/11-3 on the western margin of the Horda Platform.

*Lithology:*

The Tryggvason Formation consists generally of mudstones with interbedded limestones. Interbedded sandstones are common in the Agat area. The content of limestones relative to mudstones is generally lower in the northern part of the Viking Graben (from blocks 30/2 and 30/3 northwards) than in the southern part. At the transition between the Viking Graben and the Horda Platform (e.g. block 30/11, Fig. 36) the formation consists of limestone. The mudstones are light to dark grey, often calcareous, occasionally micaceous, glauconitic and pyritic. The limestones are white to light grey or brownishgrey and argillaceous. The sandstones are clear to light grey, very fine to fine grained and cemented by calcite.

*Basal stratotype:*

The lower boundary is defined by a decrease in gamma-ray intensity and an increase in velocity from the Blodøks Formation into the Tryggvason Formation (Fig. 33 and 34). This is due to the difference in carbonate content.

*Characteristics of the upper boundary:*

The upper boundary shows an increase in gamma-ray intensity and a decrease in velocity from the Tryggvason Formation upwards into the Kyrre Formation (Fig. 33). This log change is due to the lower carbonate content of the Kyrre Formation.

*Distribution:*

The formation is present in the Viking Graben and northern Tampen Spur area towards the Marulk Basin.

*Age:*

Early to Mid Turonian.

*Depositional environment:*

Open marine.

*Remarks:*

The Tryggvason Formation is time-equivalent with the Herring Formation and the lower part of the Hod Formation in the central North Sea, and also with the informal "formation C" of Deegan & Scull (1977) (Fig. 6).

**Kyrre Formation (new) (Kyrreformasjonen)***Name:*

Named after Olav "Kyrre" Haraldson, a Norwegian king (A.D. 1067-1093).

*Well type section:*

Norwegian well 25/1-1 from 3582 to 2997 m, coordinates N 59°53'17.40", E 02°04'42.70" (Fig. 33). Part of one core (0.5 m), including the upper boundary.

*Well reference sections:*

Norwegian well 35/3-2 from 2864 to 1665 m, coordinates N 61°51'05.98", E 03°46'28.22" (Fig. 34). No cores.

Norwegian well 24/9-1 from 3638 to 3117 m, coordinates N 59°16'09.48", E 01°41'31.18" (Fig. 35). No cores.

Norwegian well 30/11-3 from 3162 to 2892 m, coordinates N 60°02'38.59", E 02°31'15.47" (Fig. 36). No cores.

*Thickness:*

The formation is 585 m thick in the type well (25/1-1), 1199 m in well 35/3-2, 521 m in well 24/9-1 and 270 m in well 30/11-3.

*Lithology:*

The formation consists of mudstones with occasional limestone beds. Some sandstone beds are found in parts of the Agat area. The mudstones are medium grey to grey, silty to calcareous, occasionally pyritic, glauconitic or micaceous. The sandstones are clear to white, and very fine to fine grained.

*Basal stratotype:*

The lower boundary is defined by an increase in gamma-ray intensity and a decrease in velocity from the Tryggvason Formation into the Kyrre Formation (Fig. 33) due to changes in carbonate content. The boundary is unconformable on structural highs, usually above the Cromer Knoll Group.

*Characteristics of the upper boundary:*

The upper boundary shows a decrease in gamma-ray intensity and an increase in velocity from the Kyrre Formation upwards into the Jorsalfare Formation (Figs. 33 and 34). This log change is also a result of the higher carbonate content and the presence of basal limestone beds in the Jorsalfare Formation.

*Distribution:*

With the exception of the Gullfaks area, the formation is present in the Viking Graben, on the Tampen Spur and the western margin of the Horda Platform.

*Age:*

Late Turonian to Campanian.

*Depositional environment:*

Open marine.

*Remarks:*

The Kyrre Formation is time-equivalent with the Flounder Formation in the western part of the central North Sea and the upper part of the Hod Formation in the eastern part (Deegan & Scull 1977). It is also equivalent with the informal "formation D" of Deegan & Scull (1977) (Fig. 6).